Catton Grove Chalk Pit, Norwich, Norfolk

[TG 229 109]

Introduction

Catton Grove Chalk Pit (also known as 'Campling's Pit') is a small quarry surrounded by a housing estate, situated to the east of the Sprowston Road, in the Catton area of Norwich (Figure 4.24). Access is by a track from the Sprowston Road. The site provides a section across the boundary between the Weybourne Chalk and Beeston Chalk formations of the Upper Campanian Chalk of Norfolk. It is also the stratotype for the Catton Sponge Bed, which is a hardground within a complex of hardgrounds that straddles this boundary The succession here, and that at the stratigraphically higher Caistor St Edmund Chalk Pit GCR site, represents higher Campanian Chalk than is preserved in the Whitecliff, Isle of Wight and Handfast Point to Ballard Point, Dorset GCR sites. Catton Grove Chalk Pit and the nearby Attoe's Pit [TG 231 111], also now backfilled, were the source of many fossils, including ammonites, in museum collections; these came mostly from the Catton Sponge Bed and the immediately overlying beds, which are particularly fossiliferous.

The Catton Sponge Bed marks the boundary between two of the zones of the northern European Upper Campanian belemnite zonal scheme that now replaces the traditional *Belemnitella mucronata* Zone. It is a level of major macrofaunal change, which particularly affects the brachiopods, molluscs and echinoids. It marks the entry of certain benthic foraminiferal species that range up to and, in some cases, above the Campanian–Maastrichtian boundary, and is tfius of particular relevance to the interpretation of the microfaunal biostratigraphy of the Chalk successions in the southern North Sea Basin. It reflects a European-wide regressive phase, which elsewhere is expressed either by a hardground or by a change in lithofacies to shallow-water sponge-rich siliceous marls with a high-diversity macrofauna.

Description

The previously exposed section at Catton Grove Chalk Pit has been backfilled, but the Catton Sponge Bed and the overlying basal Beeston Chalk, at the top of the pit, have been deliberately covered by soil and turfed within a semi-circular retaining wall of gabions, in order to preserve it.

Lithostratigraphy

In the 1960s, this pit exposed *c*. 10 m of the highest part of the Weybourne Chalk Formation, terminating in the Catton Sponge Bed; and overlain by the basal 1.8 m of the Beeston Chalk (Peake and Hancock, 1961, fig. 6). Those workers recorded six flint bands below the Caton Sponge Bed, and one band of huge flints up to 0.45 m thick above it. They noted that an additional 22 ft (6.7 m) of section had previously been visible down to a flint that formed the floor of the pit. When this flint, probably the thick semi-tabular flint T of the stratotype Weybourne Chalk succession (Peake and Hancock, 1961, fig. 5), was broken through, the lower part of the pit rapidly flooded and had to be backfilled. Wood (1988, fig. 7) recognized some additional flints, and published a revised log. (Figure 4.26) is based on the two accounts quoted above.

The Caton Sponge Bed was named by Peake and Hancock (1961) to describe the 'hard yellow bed'or 'sponge bed' of the earlier literature that had been recorded in the chalk pits at Caxton, and which Rowe (in manuscript) and Brydone (1938) regarded as marking a level of significant faunal change in the Chalk of Norfolk. Peake and Hancock's Canon Sponge Bed comprises two closely spaced beds of iron-stained, indurated chalk capped by hardgrounds (cf. Wood, 1988, p. 62, fig. 7), rather than the single hardground recorded by them. The two hardgrounds were designated hardgrounds I and II by Wood (1988, fig. 7), the higher (the Catton Sponge Bed proper of previous workers) being taken as the boundary between his topmost (Weybourne₃) subdivision of the Weybourne Chalk and the succeeding Beeston Chalk. Hardground I is patchily and relatively poorly indurated, but is capped by a well-defined, planar erosion surface. The Canon Sponge Bed (Hardground II) contains a rich assemblage of hexactinellid sponges, in limonitic preservation, together with moulds of originally aragonite-shelled bivalves and gastropods reminiscent of the *reussianum* fauna from the Hitch Wood

Hardground of the Upper Turonian Chalk Rock. It is locally strongly indurated, and also terminates in a clearly defined erosion surface. This is overlain by soft glauconitized chalk pebbles and a concentration of large, reworked belemnites, which forms an excellent example of a so-called 'belemnite battlefield'. The Sponge Bed is penetrated by an extensive *Thalassinoides* burrow system, which in places contains belemnites 'piped down' from the overlying concentration. The huge flint A of the Peake and Hancock notation in the basal part of the Beeston Chalk is a section through a giant ring flint; it is underlain by chalk containing large pieces of shell of inoceramid bivalves, and is followed by weakly indurated chalk without an obvious erosion surface, which was designated by Wood (1988) as Hardground III.

Biostratigraphy

The section falls within the traditional *Belemnitella mucronata* macrofossil Zone, which covers the entire Upper Campanian Substage ((Figure 1.5), Chapter 1; (Figure 2.13), Chapter 2; (Figure 4.5)). There is a significant change in the belemnites, with *Belemnitella woodi* Christensen becoming extinct just below the Catton Sponge Bed, to be replaced by *B. minor I* Christensen in and above the Sponge Bed (Christensen 1995, fig. 6). The Sponge Bed marks the boundary between the *Belemnitella woodi* and *Belemnitella minor I* belemnite zones (Christensen, 1995, fig. 2; (Figure 4.5)). These zones, originally established as local zones in Norfolk, succeed the restricted *B. mucronata* Zone ((Figure 2.13), Chapter 2; (Figure 4.5)), and have now been recognized in Belgium, the Netherlands and Germany (Christensen, 1999). The belemnites in the 'belemnite battlefield' on top of the Sponge Bed, and in the basal part of the Beeston Chalk, are exclusively *Belemnitella minor* I (Christensen, 1995, fig. 6). The absence of *B. 'langel*', which is present at the base of the Caistor St Edmund Chalk Pit, places the Beeston Chalk at Canon Grove in the lowest of the three local subzones (Christensen, 1995) of the *B. minor* I Zone.

Records of ammonites (*Nostoceras (Bostrychoceras) polyplocum* (Roemer), *Baculites* sp. and *Menuites portlocki* (Sharpe)) can safely be inferred to have come from the Sponge Bed. However, other ammonites from the 'Norwich Chalk' in museum collections, particularly those preserved as glauconitized composite moulds, probably came from less well indurated ammonite-bearing horizons in the Beeston and Paramoudra Chalk. The non-ammonite molluscan fauna is largely undescribed, but includes species of the gastropod genera *Periaulax* and *Planolateralus*. The rich hexactinellid sponge fauna (details in Reid, 1968) is dominated by *Leptophragma striatopunctata* (Schrammen) with, in addition to another five species, *Aphrocallistes cylindrodactylus* Schrammen and *Lepidospongia rugosa* Schlüter. The latter two species also occur in the coeval strata in Northern Ireland.

There is a major macrofaunal change at the Sponge Bed, which especially affects the brachiopods, bivalves and echinoids. This was first noted by Rowe (in manuscript) and independently confirmed by Brydone (1922, 1938) (see review by Wood, 1988, pp. 19–39). Both workers compared this faunal change in the inland sections with the difference in faunal content between the (Weybourne) and (Beeston) Chalk successions to the west and east of Sheringham respectively. The rhynchonellid brachiopod *Cretirhynchia woodwardi* Pettitt, characteristic of the Weybourne Chalk, disappears abruptly at the top of the Sponge Bed, while the terebratulid *Carneithyris carnea* (J. Sowerby) and the rhynchonellids *Cretirhynchia arcuata* Pettitt and *C. norvicensis* Pettitt, all of which occur sporadically in the Weybourne Chalk, become abundant and represented by large-sized individuals in the Beeston Chalk. The large limacean bivalve *Plagiostoma marrotianum* (d'Orbigny) and the pectinacean *Mimachlamys mantelliana* (d'Orbigny) are apparently restricted to the pre-Sponge Bed succession. There is a striking change in the echinoids across the Weybourne Chalk–Beeston Chalk boundary, from an assemblage characterized by *Cardiotaxis heberti* Cotteau, *Micraster glyphus* Schltiter and *M. stolleyi* Lambert (the '*Epiaster'* of both Rowe and Brydone) to one characterized by *Cardiaster cordiformis* (S. Woodward) ('*Cardiaster ananchytis'*) and *Galerites roemeri* (Desor), with only extremely rare *Micraster*.

There is an important change in the microfauna (ostracods and foraminifera) across the same boundary. Two very long-ranging species of the ostracod genus *Cytherelloidea cut* out a short distance above the Sponge Bed, with other taxa entering at this level and continuing into the Maastrichtian strata (I. Slipper, pers. comm., 1998). Swiecicki (1980) recorded the extinction of some long-ranging benthic foraminiferal taxa, notably *Globorotalites micheliana* (d'Orbigny) at the Sponge Bed, and noted the abrupt entry of *Bolivina incrassata* Reuss, *Eponides beisseli* Schijfsma, *Globorotalites hiltermanni* Kaever, *Neoflabellina praereticulata* Hiltermann and *Reussella szajnochae szajnochae* (Grzybowski). These are species that range up to and, in some cases, above the Campanian–Maastrichtian boundary. There is also a significant drop in the planktonic foraminiferal content, in both numbers and diversity, at this level. The turnover in the

benthic foraminifera constitutes a significant bio-event, potentially applicable to the interpretation of offshore wells, near the top of the UKB18 or *Bolivinoides decoratus* Interval Zone (cf. Hart *et al.*, 1989, p. 314, figs 7.16, 7.25). This higher part of the UKB18 Zone, given a separate subzonal status (B3iv) by Swiecicki (1980) ((Figure 1.5), Chapter 1), is also recognized at the stratigraphically higher Caistor St Edmund Chalk Pit, and in the backfilled Frettenham Pit [TG 246 173], even higher in the Beeston Chalk. Its top is marked by the entry of *Bolivinoides miliaris* Hiltermann and Koch and *B. sidestrandensis* Barr at, or just below, the base of the Paramoudra Chalk Member in the section at West Runton confusingly labelled 'Sheringham' in the *Stratigraphical Index of Fossil Foraminifera* (Hart *et al.*, 1989, fig. 7.16).

Interpretation

South of Norwich, Hardground I and the Canon Sponge Bed were revealed in a trench in the now degraded Halfway House Chalk Pit [TG 2330 0268]. The Sponge Bed was also exposed in a trench at or near the base of the now backfilled Stoke Holy Cross Chalk Pit [TG 2536 0140] (see (Figure 4.26)). The section at the latter locality, some 9.5 km to the south along the strike from the present site, extends the succession upwards by another 2 m. In this section an additional thin, weakly indurated chalk bed and a bed of large nodular flints occur just below a weakly developed softground (Wood, 1988, fig. 7). Hardground III in the Beeston Chalk is better developed here than at Catton Grove, with a well-defined erosion surface strewn with flattened glauconitized chalk pebbles. The relationship between the composite Catton Grove–Stoke Holy Cross succession and the basal beds of the Caistor St Edmund Chalk Pit remains unclear, but it is likely that only a very small thickness of chalk separates them.

The Catton hardgrounds were formerly also seen in intermittent foreshore exposures at Sheringham, where the old Lifeboat House [TG 153 436] is actually sited on the Canon Sponge Bed. Hardground III, which can be recognized from its echinoid fauna, crops out to the east, opposite the Two Lifeboats Hotel (Peake and Hancock, 1970, p. 339E); a second hardground, possibly Hardground I, crops out a short distance to the west. The Catton Sponge Bed was not recognized in the British Geological Survey Trunch Borehole (Wood *et al.*, 1994), probably as a result of poor core recovery, but its position can be inferred from the resistivity log.

Hardground I and the Catton Sponge Bed correlate with the North Antrim Hardgrounds of Northern Ireland, which comprise two closely spaced hardgrounds, the lower one weakly, and the higher strongly, hardened and glauconitized. The higher hardground is similarly succeeded by chalks with fragmented inoceramid shell and giant ring flints (Fletcher, 1977; Fletcher and Wood, 1978). Towards the depositional margins and over structural highs, the North Antrim Hardgrounds become even more indurated and more strongly mineralized. The Catton Sponge Bed and the North Antrim Hardgrounds reflect the *polyplocum*' regression in northern Germany (Niebuhr, 1995; Niebuhr *et al.*, 1997), where it is marked by evidence of significant shallowing, including a high-diversity macrofauna with many baculitid ammonites, and the development of siliceous spongiferous marls (opoka facies), following marl-chalk rhythmites. This inter-regional regressive event, which can now be identified by correlative hardgrounds in Belgium and the Netherlands (Christensen, 1999) is interpreted as a sea-level lowstand, associated with a sequence boundary (Niebuhr *et al.*, 1997).

Conclusions

Catton Grove Chalk Pit is the type section and only remaining exposure of the Catton Sponge Bed, the other exposure on the foreshore at Sheringham having been permanently covered by the construction of a slipway. It forms the boundary between two of the belemnite zones of the standard northern European belemnite zonal scheme ((Figure 2.13), Chapter 2), and is a level of major macrofaunal change. It marks the entry of certain benthic foraminiferal species that range up to and, in some cases, above the Campanian–Maastrichtian boundary, and is thus of particular relevance to the interpretation of the microfaunal biostratigraphy of the offshore Chalk successions in the Southern North Sea Basin. It reflects a European-wide regressive phase, which elsewhere is expressed by a significant change in lithofacies to shallow water, coarse-grained, partly siliceous sediments.

References



(Figure 4.24) The location of Caistor St Edmund Chalk Pit and Catton Grove Chalk Pit, and other sections mentioned in the text, around Norwich, Norfolk. (After Cox et al., 1989.)



(Figure 4.26) The Campanian Chalk (White Chalk Subgroup) at Cation Grove Chalk Pit, Norwich, and nearby exposures (see (Figure 4.24) for locations). (Letters A-H for flint bands are those of Peake and Hancock, 1970; numbers 1–10 are those of Wood, 1988.)

Stages			fora	Benthi minif mes (I	c eral 3)	Traditional zones	Additional modern zones	Subzones		
Long			B6	ü	UKB21		Belemmella suomensis	and the second second second		
Maastrichti	ian		BS	1		Belemnella lanceolata	Belemnella obtusa			
(pars)		0		- 11	UKB20	newing much (hered)	Relevantila Lancaolata commistica			
10100000000	1		B4	i	UKB19	the state of the local	Belevenitella minor II			
	2		-	- Aler	UKBIS	Belemnitella mucronata	Belemnitella minor 1			
-	3	8	8 B3	1/10	UKB17	sensu lato	Belemnitella woodi			
a la	H	199	-	1			Beleminitella mucronata sensu structo	'Overlan mane'		
e de		10	B2	iii ii		Contraction days	committee and so being	Applications contactors		
5	wer	viec			UKB16	Gononantes dangenta	facilitative franchised franchised	прропостаная следсова		
ALL FEELS	13	s						Hagenowia blackmorei		
				- 111	1.000	Offaster pilula	15 to the second			
11/10/00			1	-			Contactional anguicas			
		1.5	B1	Same	UKB15	Marandites testudinarius	- L'arrente destructions			
=	3			1			MARK HOLD DOLLARS			
i.	den o									
mto				-		Ulmtacrimus socialis				
Sa	1.1				INPLA		None-time Associated and the			
	M		1.1		UND14		Condicerannus condiformis			
North State	L		1.1				Cladocenamus undulatoplicatus			
	U				UKB13	Michaeler corangemiem	Magadiceramus subquadratus			
	М	10		-	UKB12		Voluceramus koeneni			
acia				68	Succession in the	Micraster cortestudinarium	Inoceraneus gibbosus	These macrofossil zones are		
our	wer	10	2	TPOCT UKB11	000		Cremenoceramile created inconstant	substage concepts base		
0	3	26	19		UKB11		C. scaltersdorfensie hausonessie	largely on ammonites at		
			1			and the later of the	C. deformis erectus	inoceramid bivalves.		
	¥		3	2		francisco de altre	Prionocyclus germari	producing marker beds are		
	Upp	ini.	R zne	et al.	-	Sternozaxis plana	Subprionocyclus neptuni	also widely used (see Figure 2.3, 2.8, 2.9, 2.22 and 2.27)		
Turonian	Middle		UK Hart		UKB10	Tesebsatulina lata	Collignonicenss seocilipari			
	5		-		-	Marticles Ishina	Mammites andomides			
	5				UKB9	sensu lato	Fagesis catinus			
		-	unh			Neocardiocents inddi	Watiwocerus devonense			
	ž			14	UKB8	Metoicoceras geslinianum	and a state of the state of the			
	Up	-		13	UKB7	Galycoceras guerangeri	Constant soulies in the			
ian			1	12	UKB6	Acanthoceras jukeabrownei				
utu	fidd	(17a)	11ii 11i		UKBS	Acanthoceras		Turnilites acutus		
cuo	~	613				rhotomagense		Turrilites costatus		
0		d Hart	1	10	UKB4	Cunningtoniceras inerme Mantelliceras dixoni	The set of the below			
	ALC: N	C MD	-	9	UKB3			Mantellacensis saxbu Shantaricensis schlusteri		
	13	artes	8		UKB2	Mantellicerar mantelli		Nextline and card and		
12000	1	0	-	/	UKB1	Charles and Charles		Another the second seco		
9			-					Amaphoceras bridcensu		
bia				6		Stoliczkaia dispar		cournes antes permitations		
N			The second second				A TOTAL CONTRACTOR DE LOS CONTRACTOR	Mortonicenas (M.) rostnatum		

(Figure 1.5) Zones of the Upper Cretaceous Chalk. (* = Gap in UKB scheme; ** = UKB zonal scheme modified for this book.)

Belemnite zones NW Europe							Zonal belemnites Balto-Scandia	Zonal belemnites Russian Platform			
a dia	U		B. kauimir	viennie		28	U		20 s	U	B. kasimirostensis
SW	L	1	Br. jw	nior		Name of the	L	Top of section UK	And A	L	Br. junior
F			B. fasti	igata		F		NI and Norfolk	e		for a strain shift
-	U	-	B. cim B. sum	erceis	-	pig.	U.	~~ the		10	8. summul
- Interest		P	B. obtuss	-		Los			Lon		B. Interolata
W	1	1	l. lanceolata			W	L	B. lanceolata	X		og D. aceannin
Ug	er part	Na rones	Bt. langei	minor II	titella zone	un	icr ite	A second second	UP	υ	Bt. L majdini Bt. L langei
incom	Upp	<i>ferronite</i>	Bt. minor	minor I	Belenu	undum			undung	Ser li se la	Br. I. minor
Upper v	Lower part	itional Be	Rr.	woodi	Modern	Upper C		Bt. mucronata	Upper C	L	Bt. mucronata
		Trad	in a constraint	mucronata				B. balaskensis/Bt. mucronata	語語と		
	art Upper part	-	G. q. gracilit/B 'Overlap	t. mucronati Zone'				Bx. mammillatus G. q. scaniensis Bt. mucronata		U	Bt. mucronato/G. q. gracilig Bx. murmillatus
9			G. q. g.	ucifir					9	825	S I Street S
wer Campanu			G. q. quadrata U						wer Campaeria	м	Bt. alphaBt. praecursor/ G. q. quadesta
ŝ					ι	Lo	13				
	Lower p	G. granulataquadrata					G. granulatapuadrata Bt. alpha		L	Bt. praecurson(A, lanvigatus) G. granuloquadrata (Pteria beds)	
	υ		G, gran	wlata	U L		U	G. granulata		U	Bt. praecursor/ G. granulata
outer	-	G, sontlalica		L	G. unstialicarranulate		oniae	1	G. westfalicagramulata	1	
	М	G. westfalka			Sant	M	G. venstfalicagranulata/ Bt. propinqua/Gx. lundgreni	, j		Br. propingual	
										Gx. Insidgreni udlens	
	U		G. praces	utfalica		12121	U				
19019	M					M N	M	Gx. lundgreni	incia	U	Gx. hordgreni
3	-					- Con			Con	-	
-	L.	-	150. 1.		-	122.0	U		COLUMN STR	L	
OUIIT	M					oniae	M		onia	U	
Inc	L					1ª	L		Tor	T	E plenus triangulus
-	U	Praeactinocamax plenus				U	P. plenas	1	U	P. plenus	
e line	M		1000	1000	-	M	M		and a		
5	-		Protestimocamax primus				-	E primus	3	1	P. trimus/N. altimus
									12-12-63	*	at president to minimum

(Figure 2.13) Comparison of Upper Cretaceous belemnite zones across Europe, which are only partly represented in the UK and mainly on the Anglo-Brabant Massif. (After Christensen, 1991.) (A. = Actinocamax; B. = Belemnella; Bt. = Belemnitella; Bx. = Belemnellocamax; G. = Gonioteuthis; Gx. = Goniocamax; N. = Neohibolites; P. = Praeactinocamax.)

Stage		Southern England	Norfo	lk (Peake and Hancock, 194 Belemeites	i1, 1970) Echinoids	Norfolk (Johansen and Surlyk, 1990)	Norfolk (Christeusen, 1995, 1999)				
	Unner			Belemvella kazissirovensie							
ian	offer	Not represented Flipbere Chalk lade of Wight and Doctor Belemuitella macronata Zone Base of Zone	Not represented	Belevonitella junior	Not represented		Not represented	Belemneilla			
che			Grey Bods		Echinocorys all. limburgica	Beacon Hill Grey Chalk	NOT Y LE				
stri.			White Chalk with O. Isosata	Belemnella lichareari	Echinocorys ciplyensis	Little Marl Point Chalk Member	111111	Belemmella nomensis			
	Lower		Sponge Beds			Trimingham Sponge Beds Member-					
W			Porospharra Beds		Echisocorys belgica		Belemaitella minor II [minor III]	Belemnella obtasa			
			Sidestrand Chalk	Belenmella lanceolata	Belemmella lanceolata Echiancorys passage forma Sidestrano	Sidestrand Chalk Member		B. pseudobrasa			
			Paramata Chaite		Echisocorys pyramidata Portlock	Burnels Chill Market	Belemmitella	R. lanceolata			
a la			- and a count	Belevneitella langei dominant	1	Paramougra Chaik Member	ntinor II				
					Echinocorys conoidea Galerites roemeri-abbuesiatus	I A LINE LAN	Belevenitella minor I				
			Becitom Chaik	6.75111	Echinocorys att. conoidea Cardiotaxis ananchytis	Beeston Chalk Member					
	33		Catton Sponge Bed		Echinocorys orata auctt.	Catton Sponge Bed	101100				
nian	Upper		Wrybourne Chalk	Belemnitella macronata minor and allied forms common	Echinocorys gibba M. stolleyi	Weybourne Chalk Member	Belemmigella				
n d					Echinocorya subglobosa fonticola		woodi				
Cam			and Doeset	and Dorset	and Doeset			Echinocorya subglobosa C. beberti			
			Pre-Weybourne Chalk [Eaton Chalk]	Belemnitella mucronata activa stricto	Echinocorys pyramidata auctt, vat. quenstedi	Eaton Chalk Member	Belevonitella mocronata sense serieto				
					Echinocorys marginata approaching subgloboas						
93			Pre-Weshourne Chalk		Echinocorys lamberti						
33	12	in Hampshire	[Basal Mocronata Chalk]		Echinocorys lata fastigata	11136211					
26	Lower (park.)	Gomioteathis quadrata Zone	Gonioteuthis Zone	Gonioteathis quadrata	,						

(Figure 4.5) The 'high' Chalk of Norwich and north Norfolk based on Peake and Hancock (1961, 1970); Wood (1988); Johansen and Surlyk (1990); and Christensen (1995, 1999).