
Rhoscolyn Head RIGS site

NRW RIGS no. 242 [SH 25949 75157]

[GeoMôn Global Geopark original webpage](#)

RIGS Statement of Interest:

Rhoscolyn Head RIGS site is of regional importance as it contains the most spectacular de-watering structures on Anglesey. They have not been described elsewhere in the South Stack Formation and are important in demonstrating that the stratigraphic succession of the of this formation is the right way up. This formation consists of a half kilometer thick sequence of slightly metamorphosed sandstones deposited by a fast moving current on the sea bed (turbidites) which alternate with beds of mudstone. The de-watering structures are displayed in a bed some 70m long at the top of the cliff and can be seen from the seaward side. The water-escape structures have rounded bases which alternate with pointed crests. They represent the escape routes of unstable water-rich beds sandwiched between more coherent layers of mudstone. These rare features are the only known examples in the Precambrian rocks of Anglesey and help us to understand the conditions and environment of deposition at a time around 500–600 million years ago. One possible explanation for their existence is that the unit marks a time of instability resulting from an earthquake.

Geological setting/context: The Precambrian basement rocks of Anglesey and south-west LI■n can be divided into several discrete groups, all of which were juxtaposed along a series of steep, brittle and/or ductile faults and shear zones (e.g. Dinorwic and Aber-Dinlle faults; Berw, Central Anglesey and LI■n shear zones) collectively referred to as the Menai Strait Fault System (MSFS). First, the Monian Supergroup consists of a thick sequence of polydeformed metasediments and meta-igneous rocks, comprising the South Stack, New Harbour and Gwna groups, the latter representing the type example of a large-scale submarine debris flow or *mélange* said by some researchers to be of Lower Cambrian age. Ongoing research, however, may suggest a much older date for the Gwna Group with possible Cambrian ages being put forward for the South Stack metasediments. Second, the Coedana Complex of central Anglesey comprises high-grade metasediments, amphibolites and gneisses, and low-grade, thermally metamorphosed hornfeldes adjacent to a granite (Coedana Granite), which has recently yielded a late Precambrian zircon age of $614 \pm 4\text{Ma}$. Third, a belt of schists and metabasites displaying blueschist facies grade of metamorphism lies within the MSFS. The metabasites exhibit a strong mid-ocean ridge basalt signature and have yielded ages of 580–590Ma. Fourth, the Sarn Complex in LI■n comprises metagabbros and granite rocks which occur to the south-east of the LI■n Shear Zone (LSZ), a continuation of the MSFS, which separates these igneous rocks from low-grade Monian *mélange* to the north-west. A late Precambrian zircon magmatic age of $615 \pm 2\text{Ma}$ has been obtained from a metagabbro of the LSZ. Fifth, on the mainland of north-west Wales, the Arfon Group comprises a thick sequence of tuffs and volcanoclastic rocks, dated at $614 \pm 2\text{Ma}$, which are conformably overlain by late Lower Cambrian siltstones. Correlatives of the Arfon Group may occur as isolated outliers on Anglesey and, if proven, would provide an important potential lithostratigraphical link across the MSFS. The stratigraphical correlation between the various units has proved highly controversial.

The recent recognition of mylonitic rocks, for example in the LSZ, emphasises the presence of tectonic contacts and indicates that each component may represent a so-called 'suspect terrane' which was transported laterally into position along the major faults and shear zones. Ongoing unpublished research suggests, that Anglesey's Precambrian rocks accumulated in accretionary prisms, providing a tectonic sequence rather than a stratigraphic sequence which was formerly accepted. This new research would reverse the accepted stratigraphic order established for the island. This Precambrian basement later formed the north-west margin of the Lower Palaeozoic Basin, the initiation of which was contemporaneous with Arfon Group volcanism. The timing of the inferred fault displacements has also been the subject of debate. Investigations on LI■n have demonstrated that assembly of the basement terranes was completed at least by early Ordovician times since an unconformable Arenig overstep sequence has been identified at several localities such as Wig Bach, Parwyd and Mountain Cottage Quarry. The Arenig sequence of Anglesey and LI■n is considerably less deformed and metamorphosed than the underlying basement, although this distinction is not everywhere obvious.

To select RIGS to demonstrate the Precambrian evolution of Anglesey and Llŷn, three separate networks were devised. These are: 1. Precambrian stratigraphy and structures. This network includes two sub-sets: a) Precambrian sedimentary structures; and b) tectonic structures, such as folds and faults, which may have occurred during a tectonic event in Precambrian times or later, for example, during the Caledonian Orogeny; 2. Precambrian palaeontology which includes any life-form and trace fossil, such as stromatolites, sponge spicules, worm burrows and bioturbated metasediments. Current research suggests that some of these fossils may be Cambrian or even Ordovician in age, but as these life-forms were previously held to be Precambrian in age, they have been included in this category; and 3. Precambrian reference sections. These aim to represent all important Precambrian rock types found in Anglesey and Llŷn. They include the major mapped units of Greenly (1920). The aim is to provide the best and most accessible exposure of the rock type. These can be considered as RIGS 'type sections'. Where there is a relevant mineralogical, sedimentary, structural or other change across an outcrop, several representative sites have been chosen.

Network context of the site: Rhoscolyn Head is designated for the important sedimentary structures, namely, de-watering features, contained in a bed some 70m long at the top of the steep sea cliffs within the South Stack Formation. It is therefore designated category 1a. This site is the sole sedimentary structure to represent this category within the Precambrian Structures Network.

References:

GREENLY, E. (1919). The geology of Anglesey. Memoirs of the Geological Survey of Great Britain. HMSO, London, 980pp. (2 vols)

GREENLY, E. (1920). 1:50,000 (and 1 inch to 1 mile) Geological Map of Anglesey. Geological Survey of Great Britain, Special Sheet No. 92 and (93 with parts of 94, 105 and 106).

SHACKLETON R. M. (1975). Precambrian rocks of North Wales . In WOOD, A, (ed). The Pre-Cambrian and Lower Palaeozoic rocks of Wales. University of Wales Press, Cardiff, 1–22.

WOOD, M. (2003). Precambrian Rocks of the Rhoscolyn Anticline. Seabury Salmon & Associates, Ludlow, 1–48.